

Clinical Refraction, Accommodation, Refractive Errors, and Astigmatism

Mardanov Botirbek Akmalovich

Student, Group D1-404A, Faculty of General Medicine,
Termez Branch of Tashkent State Medical University,
Uzbekistan mardanovbotirbek@gmail.com
<https://orcid.org/0009-0000-6933-8948>

Esonova Nafisa Alisherovna

Assistant, Department of Otorhinolaryngology,
Ophthalmology, Oncology and Medical Radiology,
Termez Branch of Tashkent State Medical University ,Uzbekistan
nafisaesanova1989@gmail.com
<https://orcid.org/0009-0006-4524-5302>

Abstract: Clinical refraction is a fundamental component of ophthalmic practice and optometric assessment because it provides an objective and subjective evaluation of the optical status of the eye and determines the appropriate refractive correction. Clear vision depends on the precise interaction between the cornea, crystalline lens, aqueous and vitreous media, retina, and neural visual pathways. Among these structures, the cornea and crystalline lens provide most of the eye's refractive power, while accommodation enables the eye to maintain clear retinal images when viewing objects at different distances. Disturbances in the optical system result in refractive errors, including myopia, hyperopia, and astigmatism. Astigmatism is particularly important because it produces meridional differences in refractive power and may significantly affect visual acuity and visual quality. This article reviews the principles of clinical refraction, physiological mechanisms of accommodation, major types of refractive errors, and the clinical characteristics, diagnosis, and correction of astigmatism. Particular attention is given to modern approaches to refractive assessment and the importance of individualized optical correction.

Keywords: clinical refraction, accommodation, refractive error, myopia, hyperopia, astigmatism, visual acuity, optometry, ophthalmology.

Introduction

For normal distance vision, parallel rays of light entering the eye must be appropriately refracted so that they converge on the retinal photoreceptor layer. The refractive status of the eye is determined primarily by the optical power of the cornea and crystalline lens, as well as by the axial length of the globe and the refractive indices of the ocular media. When the optical system does not focus light precisely on the retina in the absence of accommodation, a refractive error is present. Refractive errors are among the most common causes of reduced visual acuity worldwide. The principal forms are myopia, hyperopia, and astigmatism. Presbyopia represents an age-related reduction in accommodative ability rather than a classical refractive error, although it has major clinical importance in refractive practice. Clinical refraction is the process by which the refractive

status of an individual is assessed and an appropriate optical correction is determined. It includes objective techniques, such as retinoscopy and automated refraction, and subjective techniques in which the patient compares different lens powers while reporting perceived visual clarity. Accurate refraction requires an understanding of ocular physiology, accommodation, binocular vision, visual acuity, and the limitations of different measurement methods.

MATERIALS AND METHODS

This article was prepared as a descriptive review of basic ophthalmological concepts related to clinical refraction, accommodation, refractive errors, and astigmatism. The analysis focused on the physiological and optical principles of the eye, common types of refractive abnormalities, methods of clinical assessment, and approaches to optical correction. The main areas considered in the review included:

1. **Clinical refraction** — principles and methods used to determine the refractive status of the eye.
2. **Accommodation** — physiological mechanisms responsible for maintaining clear vision at different distances.
3. **Refractive errors** — characteristics of myopia, hyperopia, and age-related changes in accommodation.
4. **Astigmatism** — its optical characteristics, clinical manifestations, diagnosis, and correction.

Both objective and subjective approaches to refractive assessment were considered, with attention to the importance of individualized correction based on visual acuity and patient needs.

RESULTS

Clinical refraction involves determining the optical power required to produce a clear retinal image. During the examination, visual acuity is assessed and the refractive state of each eye is evaluated. Objective refraction can be performed using retinoscopy or an autorefractor. These methods provide an initial estimation of the refractive error without requiring the patient to select the lens power. Subjective refraction is performed by presenting different spherical and cylindrical lens powers and asking the patient to compare visual clarity. This process allows the examiner to refine the optical correction according to the patient's visual response. Accurate refraction requires consideration of accommodation, binocular vision, pupil size, visual demands, and the patient's age. In some patients, excessive accommodation may influence the measured refractive state, particularly during near or subjective testing. Accommodation is the physiological ability of the eye to increase its optical power in order to maintain a focused image of a near object on the retina. The crystalline lens plays the central role in this process. When viewing a distant object, the ciliary muscle is relatively relaxed and the lens has a flatter shape. During accommodation for near vision, contraction of the ciliary muscle reduces tension on the zonular fibers, allowing the crystalline lens to become more convex. This increases its refractive power and permits the eye to focus on nearby objects. Accommodation is associated with the near response, which also includes convergence of the eyes and pupillary constriction. These mechanisms work together to provide clear and comfortable near vision. With increasing age, the crystalline lens gradually loses its ability to change shape effectively. This age-related reduction in accommodation is known as presbyopia and commonly results in difficulty focusing on near objects. A refractive error occurs when the optical system of the eye does not focus light precisely on the retina when accommodation is relaxed. The main refractive errors are myopia, hyperopia, and astigmatism. **Myopia** occurs when parallel light rays are focused in front of the retina when accommodation is relaxed. This may result from excessive axial length of the eye, excessive refractive power, or a combination of these factors. Patients with myopia generally experience reduced distance visual acuity while near vision may remain relatively clear. **Hyperopia** occurs when the optical system would focus parallel light rays behind the retina in the absence of accommodation. It may be associated with a shorter axial length

of the eye or insufficient optical power. Depending on its degree and the patient's accommodative ability, hyperopia may cause blurred vision, visual fatigue, or difficulty with near tasks. **Presbyopia** is an age-related reduction in accommodative ability. It differs from myopia and hyperopia because the primary problem is the loss of accommodation rather than an abnormal static refractive state. Patients commonly report difficulty reading small print or performing prolonged near work. Astigmatism is a refractive condition in which the optical power differs between different meridians of the eye. Instead of forming a single point focus on the retina, incoming light may form different focal lines or planes. As a result, the patient may experience blurred, distorted, or reduced-quality vision. Astigmatism may be classified according to the orientation and relationship of the principal meridians. Regular astigmatism occurs when the principal meridians are approximately perpendicular to one another, whereas irregular astigmatism may result from an uneven or irregular optical surface. Astigmatism can be associated with the cornea, the crystalline lens, or both. Corneal astigmatism is particularly common because small differences in corneal curvature between meridians can significantly affect the refractive power of the eye. Clinical symptoms of astigmatism may include blurred vision at different distances, visual fatigue, headaches, difficulty with prolonged visual tasks, and distortion of objects. The severity of symptoms depends on the magnitude and type of astigmatism as well as individual visual requirements.

DISCUSSION

Clinical refraction is an important part of comprehensive eye examination because refractive abnormalities can significantly affect visual performance and quality of life. Objective measurements provide useful information about the optical state of the eye, while subjective refraction allows the prescription to be adjusted according to the patient's visual perception. Accommodation is particularly important when assessing refractive status. Changes in accommodative activity can influence refractive measurements, especially in younger patients with strong accommodative responses. Therefore, appropriate examination techniques and careful interpretation of results are necessary to obtain reliable measurements. Myopia and hyperopia primarily involve differences between the optical power of the eye and its axial length, whereas astigmatism involves differences in refractive power between meridians. Although these conditions have different optical mechanisms, all may lead to reduced visual acuity if they are not appropriately corrected. Astigmatism deserves particular attention because it may cause both blurred and distorted vision. Cylindrical correction can compensate for differences in refractive power between the principal meridians. In cases involving irregular corneal surfaces, additional diagnostic procedures may be necessary to determine the most appropriate management. The selection of refractive correction should therefore be individualized. A prescription should not be based solely on an automated measurement. The clinical findings, subjective responses, visual requirements, and overall ocular condition should be considered together.

CONCLUSION

Clinical refraction is a fundamental component of ophthalmic assessment and plays an important role in identifying and correcting refractive abnormalities. Accommodation enables the eye to maintain clear vision at different distances through changes in the optical power of the crystalline lens. Myopia, hyperopia, and astigmatism represent major refractive conditions that may reduce visual acuity and affect visual performance. Astigmatism is characterized by different refractive powers in different meridians and may produce blurred or distorted vision. Accurate diagnosis requires appropriate objective and subjective refractive assessment, while individualized optical correction can improve visual acuity and visual comfort. A thorough understanding of clinical refraction, accommodation, and refractive errors is essential for medical students, ophthalmologists, and optometrists. Continued development of objective diagnostic technologies and individualized approaches to refractive correction may further improve the assessment and management of patients with refractive abnormalities.

REFERENCES:

1. Benjamin W.J. **Borish's Clinical Refraction**. 2nd ed. St. Louis: Butterworth-Heinemann; 2006.
2. Elkington A.R., Frank H.J., Greaney M.J. **Clinical Optics**. 3rd ed. Oxford: Blackwell Science; 1999.
3. Kanski J.J., Bowling B. **Clinical Ophthalmology: A Systematic Approach**. 8th ed. Elsevier; 2016.
4. American Academy of Ophthalmology. **Basic and Clinical Science Course: Optics and Refraction**. San Francisco: American Academy of Ophthalmology.
5. Atchison D.A., Smith G. **Optics of the Human Eye**. Oxford: Butterworth-Heinemann; 2000.
6. Grosvenor T. **Primary Care Optometry**. 5th ed. St. Louis: Butterworth-Heinemann; 2007.
7. Elliott D.B. **Clinical Procedures in Primary Eye Care**. 4th ed. Elsevier; 2014.
8. Yanoff M., Duker J.S. **Ophthalmology**. 5th ed. Elsevier; 2019.