

TEST METHODS USED TO DETECT BREAST CANCER IN THE EARLY STAGES

Ruziyeva Shahlo Durdiqulovna

Bukhara State Medical Institute

Relevance:

When breast cancer is suspected, smearing, palpation, examination of both mammary glands by ultrasound and mammography methods, cytological examination of the punctate obtained from the tumor, and histological examinations with sectoral resection, if necessary, are carried out. Examination is of great importance for diagnosis. Factors that cause cancer can be determined by the presence of diseases in the genital organs from a young age, the time when menstruation starts and stops, the number of pregnancies and abortions, breastfeeding, the use of hormonal drugs, and other questions. Mammography is performed using special x-ray diagnostic equipment without any contrast agents. The method of fluoromammography is rarely used in recent years. X-rays are taken in bottom and side projections. Usually, mammography is performed in such projections for comparison. Examination of young women is performed between menstruation. In mastopathies, it is possible to see diffuse and fibrous changes in the structure of the mammary gland, and between them small lights (cysts) have appeared. While the mammography method has a high level of diagnostic value in the examination of the mammary gland, one should not forget its carcinogenicity (cancer formation in the gland). Therefore, this method can be used only 1-2 times in young people. This method should be used mainly in women over 50 years of age. Ultrasound examination method (sonography). In recent years, this method has been widely used in the examination of mammary glands. It clearly identifies cystic and nodular changes. Since this method is harmless to the mammary gland, it can also be used in young patients. Other special examinations (thermography, ductography, transillumination, xeromammography, contrast mammography, arteriography, radioisotope diagnostics) are not widely used for some reasons. Radioisotope examination is mainly used to detect bone metastases. When changes in the mammary gland are examined by all available methods, the diagnosis of cancer is made only after pathomorphological, cytological and histological examinations. For cytological examination, the skin of the mammary gland is pierced with a special needle, and the contents of the nodule or lesion are drawn into a syringe (puncture biopsy) and microscopic examination is performed. In this case, abnormal (atypical) and cancerous cells can be found in the mammary gland.

Objective: To provide clinical and morphological justification for modern complex treatment methods for breast cancer.

Materials and Methods: Study Types and Objects: 150 patients with breast cancer treated between 2018 and 2023 were clinically and morphologically studied. Patients were divided into groups according to their molecular type and stage of disease

Result, Two semi-circumcised skin incisions are made in the radial direction, 2 cm away from the border of the tumor. One of the incisions goes deep to the pectoralis major muscle. The left ash fingers are inserted into the resulting bush, and the sector to be removed is waiting.

Conclusion, an incision on the opposite side of the carapace with connective tissue is deepened, and the damaged sector is cut. In a technically correct resection, the bottom of the wound is the fascia of the pectoral muscle. The removed tumor is surrounded by mammary tissue from all sides. When nodular mastopathy, fibroadenoma and cancer are suspected, sectoral resection is performed in the same way, without any technical difference.

Referances:

1. Беспалов В.Г. Лечение мастопатии и первичная профилактика рака молочной железы. Лечащий врач 2017;(5):88-9.

2. Васильев Д.А., Зайцев А.Н., Берштейн Л.М. Маммографическая плотность молочных желез и определяющие ее факторы в свете повышенного онкологического риска. Опухоли женской репродуктивной системы 2011;(3):15-22.

3. Высоцкая И.В., Погодина Е.М., Гладилина И.А. и др. Клиническая маммология (практическое руководство). Под ред. М.И. Давыдова, В.П. Летягина. /М., 2010;54-6.

4. Герштейн Е.С. Биологические маркеры молочной железы: методологические аспекты и клинические рекомендации /Е.С. Герштейн, Н.Е. Кушлинский //Маммология.2015;1:65 69.

5. Каприна А.Д. и др. Злокачественные новообразования в России в 2013 году (заболеваемость и смертность). Под ред. А.Д. Каприна, В.В. Старинского, Г.В. Петровой. М., 2015;250.

6. Зотов А.С., Белик Е.О. Мастопатии и рак молочной железы. /М., 2005;112. Высоцкая И.В. [и др.] Опухолевые маркеры рака молочной железы //Маммология. 2015;1:61-65.

7. Канцерогенез. Под ред. Д. Г; Заридзе. М., 2005 Коган И.Ю., Мясникова М.О. Диагностика и лечение мастопатии. СПб., 2010;32.

8. Летягин В.П. Факторы риска развития рака молочной железы / В.П. Летягин, И.В. Высоцкая, Е.А. Ким // Маммология. 2006;4:10-13.