

UDC:616.38-002

CHARACTERISTICS OF THE PULMONARY ENDOTHELIAL SYSTEM IN ABDOMINAL SEPSIS

Khotamov Ihtiyor Eshmurotovich, Khamdamov Bakhtiyor Zarifovich
Bukhara State Medical Institute. Bukhara, Uzbekistan.

Summary. Molecular-biochemical parameters of the endothelial system of the lungs during experimental modeling of abdominal sepsis showed an ambiguous picture of the changes occurring, which were of a phase nature and depended on the timing of the development of the pathological process.

Key words: Abdominal sepsis, lungs, endothelial system.

Relevance. Throughout the last century, abdominal sepsis has been and remains a dangerous disease in which dysfunction of vital organs develops as a result of the aggression of introduced microorganisms.

The high incidence of abdominal sepsis is accompanied by high mortality, which can vary from 7.6% to 36.0% (1,3,5,25,26,27,28,29,30,31,32,33). Intra-abdominal contamination and secondary peritonitis are a constant source of pathogen-associated molecular patterns (via spillage of intestinal contents) and through direct damage to internal and abdominal organs. This “engine of multisystem organ failure” provides continuous cytokine fuel for the raging systemic response (2,4,6,8). For example, TNF- α and IL-1 are important proinflammatory cytokines. Each has been shown to induce vascular permeability, leading to pulmonary edema and hemorrhage (7,9,11). IL-6 is a key molecule in initiating the febrile response, lymphocyte activation, and also plays a role in hematopoiesis. It has also been shown to cause myocardial depression (10,12,14,16,18).

Over the past 20 years, the recognition of the endothelium as a full-fledged system has led to a large number of experimental and clinical studies, including the study of the mechanisms of development of sepsis. During bacterial, fungal, or viral infection, exogenous molecular patterns associated with pathogens and molecular patterns associated with endogenous damage cause endothelial activation and can disrupt its structure and function, that is, provoke the development of endothelial dysfunction (13,15,17,19,34,35,36,37,38,39). Endothelial changes associated with sepsis should be considered useful to limit the spread of bacteria, as well as to control leukocyte recruitment and bacterial clearance. However, severe and persistent phenotypic changes in the endothelium may contribute to impaired microcirculatory blood flow, tissue hypoperfusion, and the development of life-

threatening multiple organ failure (21,23,24,40,41). Thus, it is very difficult to distinguish between corresponding activation and endothelial dysfunction, especially given that the endothelial cell response may vary between organs (20,22). Purpose of the study: to study the endothelial system of the lungs in abdominal sepsis. Material and methods. The studies were conducted on laboratory outbred rats in a model of abdominal sepsis that we developed. The experiments were carried out on 106 white Wistar laboratory rats weighing 200-250 grams, of both sexes, fed a regular laboratory diet. The planned experimental studies, which included sampling, biopsies and necropsies, were based on the conditions specified in the 1986 Council of Europe Animal Welfare Convention. The animals were divided into the following series of experiments: Control – 10 intact animals, not subjected to any influences or manipulations, fed a standard grain diet. The main one was 50 animals in which the experimental model of abdominal sepsis was reproduced using our improved method. The reproduction of the experimental model of abdominal sepsis was carried out in stages, by changing the reactivity of animals and creating a purulent-necrotic focus in the abdominal cavity. In experimental studies, blood was collected separately at the entrance and exit from the lungs. In this case, the blood at the entrance to the lungs is mixed venous blood, which came from the inferior and superior vena cava. At the exit from the lungs we received arterial blood, which was universal for the entire organism as a whole. This technique was developed and tested by a group of researchers at the Tashkent Medical Academy. Each value obtained in different blood samples was also subjected to calculation of the venous-arterial difference, that is, a value reflecting the “delay” or “production” of the substrate in the endothelial system of the lungs.

The entire complex of studies of patients with abdominal sepsis consisted of continuous monitoring of the state of homeostasis and functional activity of vital organs. For this purpose, functional, instrumental and laboratory research methods were carried out. Integral rating diagnostic methods, such as APACHE II, SAPS, SOFA, and the Kalf-Kalif Leukocyte Intoxication Index, were also actively used. Among the indicators of the endothelial system, the level of C-reactive protein (mg/l), thrombomodulin (ng/ml), von Willebrand factor (IU/dL), intercellular and cellular adhesion molecules (ng/ml) was studied using an enzyme-linked immunosorbent analyzer; nitrites and nitrates (%), peroxynitrite ($\mu\text{mol/l}$), nitric oxide synthase activity ($\mu\text{mol/min/l}$) according to the Griess method modified by A.P. Solodko et al. on an SF-46 spectrophotometer at a wavelength of 520 nm. The entire set of studies met the criteria of translational medicine, which used the entire body of research, which

made it possible to extrapolate the results of experimental studies into clinical practice. Results and its discussion. The study of the total value of NO is considered a generally accepted indicator of the nitroxergic system for regulating vascular tone, as one of the functional criteria of the endothelial system. The average level of its content in the mixed venous blood sample at the entrance to the lungs was $26.39 \pm 3.91 \mu\text{mol/l}$, while at the exit from the lungs in the arterial blood sample its level was higher and reached an average value of $33.41 \pm 4.64 \mu\text{mol/l}$. The venous-arterial difference, which averaged “+” $7.02 \pm 1.13 \mu\text{mol/l}$, was positive and indicated the production of this element in the alveolar capillary network with release into the systemic bloodstream. It should be noted that this nature of the venous-arterial difference was typical in all series of experiments we studied. In this case, the peak values occurred in animals of the control and comparative groups (“+” $8.44 \pm 2.17 \mu\text{mol/l}$ and “+” $8.63 \pm 2.35 \mu\text{mol/l}$, respectively). In other cases, in the dynamics of development of the experimental model of abdominal sepsis, the production of this substrate of the nitroxergic system for regulating vascular tone decreased from “+” $8.11 \pm 2.47 \mu\text{mol/l}$ ($p < 0.05$) for the 6-hour period of reproduction of the experimental model of abdominal sepsis and up to “+” $4.69 \pm 1.62 \mu\text{mol/l}$ ($p < 0.05$) for a 24-hour period of disease development. In the subsequent 48, 72 and 96-hour periods of development of the experimental model of abdominal sepsis, the venous-arterial difference in NO increased again. However, the venous-arterial difference did not reach the initial value. Moreover, most of the indicators were of unreliable changes, reflecting the distance from the ongoing real processes associated with the development of the experimental model of abdominal sepsis. In this regard, we conducted a dispersion analysis of the constituent elements of NO (NO₂⁻ and NO₃⁻) in percentage terms in the dynamics of the development of an experimental model of abdominal sepsis. The predominant role in the percentage value of NO₂⁻, already for a 24-hour period of modeling abdominal sepsis, is leveled out, giving way to the fractional value of NO₃⁻. In other words, there is an increased production of NO₃⁻ by the endothelial system in the conditions of ongoing disorders. This once again confirms the importance of assessing the component of NO decay rather than its overall value. Against the background of the changes described above, we identified changes in the concentration of the metabolic product of the transformation of NO components, in particular NO₃⁻ into OONO⁻. The average content of peroxynitrite in the mixed venous blood sample at the entrance to the lungs throughout the study exceeded its level in the

arterial blood sample at the exit from the lungs ($2.5 \pm 0.09 \mu\text{mol/L}$ and $2.37 \pm 0.03 \mu\text{mol/L}$; $p > 0.05$).

The venous arterial difference, which was negative (“-”), indicated the active utilization of this oxidative product in the endothelial system of the lungs and a decrease in its production into the systemic arterial bloodstream. In other words, the lungs, and in this case its endothelial system, performed a barrier filtration function, creating conditions for the formation of blood that was universal in composition for all organs of the body. We identified minimal values with an unreliable level of differentiation between the content of peroxynitrite in the mixed venous blood sample at the entrance to the lungs and in the arterial blood sample at the exit from the lungs among animals in the control and comparative series of experiments. The venous-arterial difference (“-” $0.03 \pm 0.01 \mu\text{mol/l}$), equivalent to 10%, can be safely taken as absent. Meanwhile, starting from the 24-hour period of development of the experimental model of abdominal sepsis, there was a decline in the venous-arterial difference to “-” $0.74 \pm 0.12 \mu\text{mol/l}$ ($p < 0.05$), which in subsequent periods led to an inversion of values, due to a change in the nature of the formation of peroxynitrite, which began to be actively synthesized (“+”) in the endothelial system of the lungs. Its increase in the arterial blood sample at the exit from the lungs, starting from this period of development of the experimental model of abdominal sepsis, was significantly pronounced, especially at 48 hours (7.4 times relative to 6 hours and 7.1 times relative to 12-hour periods) and to 72-hour (8.9 times relative to 6-hour and 8.5 times relative to 12-hour periods) periods. Thus, it is possible to state an increase in peroxynitrite in various blood samples depending on the timing of the development of the experimental model of abdominal sepsis, which was characterized by a transition from a state of transient phenomena to steadily progressive ones, indicating the depletion of the compensatory capabilities of the pulmonary endothelial system itself. As evidence for the above conclusion, we consider the dynamics of changes in the activity of the iNOS enzyme in the studied blood samples at the entrance and exit from the lungs. In particular, in intact animals the activity of this enzyme is inhibited as it passes through the endothelial system of the lungs. The same character of the ratio of endothelial cells can be seen in the ratio of the group of animals of the comparative series. It is interesting that the venous-arterial difference continues to increase among animals with an experimental model of abdominal sepsis. However, as in the case of peroxynitrite, this pattern completely changes starting from the 24-hour period of development of the experimental model of abdominal sepsis. In the

dynamics of development of the experimental model of abdominal sepsis, identical changes occurred in the concentration of intercellular and cellular adhesion molecules. In intact animals, a decrease in the concentration of these molecules was observed as blood passed through the endothelial system of the lungs. The venous-arterial content of ICAM-1 decreased by 29.8 times ($p < 0.001$ - significant value in the arterial blood sample at the exit from the lungs in relation to the mixed venous blood sample at the entrance to the lungs).

In terms of VCAM-1 content, the decrease was slightly less - 22.9 times, although it was also stable and significant ($p < 0.001$ - significant value in the arterial blood sample at the exit from the lungs in relation to the mixed venous blood sample at the entrance to the lungs). Such changes correspond to physiological parameters, which are confirmed by the presence of high tone in the arterial circulatory system. Also interesting is the identical level of change in the venous-arterial difference (24.4 times, respectively) for both indicators. This occurred due to a decrease in the specific gravity of intercellular adhesion molecules and an increase in cell adhesion molecules in the mixed venous blood sample at the entrance to the lungs. When modeling abdominal sepsis, starting from the 6-hour study period, we noted a progressive decrease in the level of venous-arterial difference. At the same time, the minimum value in relation to the venous-arterial difference of ICAM-1 was noted for the 24-hour period of development of the experimental model of abdominal sepsis, while in relation to VCAM-1 - for the 48-hour period of development of the experimental model of abdominal sepsis. This was apparently due to the stages of changes occurring in the endothelial system of the lungs, where at the first stage there was a predominance for intercellular adhesion, and subsequently - cellular adhesion, which indicates the presence of endothelial apoptosis. We also noted a leveling of the values of the venous-arterial difference in this period of experiments in relation to vWF. The maximum level of vWF formation in the endothelial system of the lungs occurred during the 12-hour period of development of the experimental model of abdominal sepsis (1.2 times). In subsequent periods, as the pathological process progressed, an increase in the venous-arterial difference over the 24-48-hour study period indicated a cumulative characteristic between vWF in the venous blood sample at the entrance to the lungs and in the arterial blood sample at the exit from the lungs. This nature of the changes was reflected in the subsequent development of the experimental model of abdominal sepsis, in which the productivity of vWF in the endothelial blood system at the exit from the lungs indicated active structural and functional disorders already in the pulmonary vessels. Analysis of the venous-arterial difference showed

that the endothelial system of the lungs reacted sensitively to changes occurring in the site of destruction. At the same time, the main character of the endothelial system of the lungs at the first stage was reduced to blocking the flow of pathological substrates into the systemic arterial bloodstream, and at the second stage of development of the experimental model of abdominal sepsis, the lungs cease to create a barrier to the generalization of the inflammatory process, opening the way for the development of multiple organ dysfunction.

Conclusions. 1. The increase in peroxynitrite in various blood samples depending on the timing of development of the experimental model of abdominal sepsis, which was characterized by a transition from a state of transient phenomena to steadily progressive ones, indicating the depletion of the compensatory capabilities of the pulmonary endothelial system itself.

2. The productivity of vWF in the endothelial blood system at the exit from the lungs was evidenced by active structural and functional disorders already in the pulmonary vessels.

Literature

1. Adihamzhanov Sh. B., Muminov B. E. Dates of initiation and volumes of nutritional support for abdominal sepsis // *Economics and society*. – 2022. – No. 3-2 (94). – pp. 407-417.
2. Afonasyeva T.M. Endothelial dysfunction. Availability of early diagnosis. *Health and education in the 21st century*. 2016;18(11):101–4.
3. Babadzhanov B.D., Okhunov A.O., Atakov S.S. Methodology for studying the metabolic function of the lungs in small laboratory animals // *Methodological recommendation*. – 1992.-Tashkent, 34 p.
4. Barkhatova N. A. Peritonitis and abdominal sepsis: etiology, systemic response, choice of antibacterial therapy // *Indelible tablets: sepsis et cetera: Collection of materials from the conference of the Association of General Surgeons, dedicated to the anniversary of the Department of General Surgery of Yaroslavl State Medical University, Yaroslavl, May 18–19 2020*. – Yaroslavl: Digital Printing House, 2020. – pp. 41-45.
5. Clinical and nosological characteristics of peritonitis and abdominal sepsis in a city multidisciplinary hospital during the COVID-19 pandemic / A. A. Naidenov, A. V. Gostimsky, I. V. Karpatsky, M. V. Gavshchuk // *Journal of Emergency Surgery named after . I.I. Dzhanelidze*. – 2021. – No. S2. – pp. 53-54.
6. Determination of the optimal fluid balance in patients with abdominal sepsis in the postoperative period / S. A. Tochilo, A. V. Marochkov, T. I. Klepcha [et al.] // *Surgery of Belarus - state and development: collection of scientific materials*

practical conference with international participation and the XVII Congress of Surgeons of the Republic of Belarus, Mogilev, October 12–13, 2023. – Minsk: Belarusian State Medical University, 2023. – P. 546-547.

7. Ostroumova Yu. S. Comparative assessment of criteria for predicting the course of secondary peritonitis and abdominal sepsis: specialty 01/14/17 “Surgery”: dissertation for the degree of candidate of medical sciences / Ostroumova Yulia Sergeevna, 2021. – 110 p.

8. Khamdamov I.B. Improving tactical approaches in the treatment of hernias of the anterior abdominal wall in women of fertile age // New day in medicine. Bukhara, 2022.-№10(48)- P. 338-342.

9. Khamdamov I.B. Morphofunctional features of the abdominal press in women of reproductive age // New day in medicine. Bukhara, 2022.-№3(41)- P. 223-227.

10. Khamdamov B.Z. Comparative evaluation of methods of amputation related to tidiotarus with severe forms of diadetic foot syndrome. European Science Review. Austria, Vienna 2014 Septemba-October No. 9-10. - pp. 58-60.

11. Khamdamov B.Z. Diabetic tovon syndromida bazhariladigan yukori amputationlardan sung bemorlarning hayot kechirish sifatidagi uzgarishlar takhlili. Samarkand. Biology va tibbiyot muammolari. No. 1, 2019., (107) - pp. 115-117.

12. Khamdamov B.Z. Complex treatment of diabetic foot syndrome with critical ischemia of the lower extremities. Journal of Biomedicine and Practice. Tashkent 2020, Special issue. Part 5 – pp. 801-814.

13. Khamdamov B.Z. Method of laser photodynamic therapy in the treatment of wound infection in diabetic foot syndrome. Biology va tibbiyot muammolari No. 1 (116) 2020. – P. 142-148

14. Khamdamov B.Z. Morphological changes when using photodynamic therapy in the treatment of wound infection in an experiment. Journal of Morphology. Saint Petersburg. 2020. Volume 157 (2-3). -WITH. 223-224.

15. Khamdamov B.Z. Optimization of methods for local treatment of purulent-necrotic lesions of the foot in diabetes mellitus. Magazine. Tibbiyotda yangi kun. 2018, No. 4 (24) - pp. 112-115.

16. Khamdamova M.T. Ultrasound features of three-dimensional echography in assessing the condition of the endometrium and uterine cavity in women of the first period of middle age using intrauterine contraceptives // Biology va tibbiyot muammolari. - Samarkand, 2020. - No. 2 (118). - P.127-131.

17. Khamdamova M. T. Ultrasound assessment of changes in the endometrium of the uterus in women of the first and second period of middle age when using

- intrauterine and oral contraceptives // Biomedicine va amaliyot journals. – Tashkent, 2020. - No. 2. - Part 8.- P.79-85.
18. Khamdamova M. T. Anthropometric characteristics of the physical status of women in the first and second period of middle age // A new day in medicine. Tashkent, 2020. - № 1 (29). - P.98-100.
19. Kaur H, Chandran VP, Rashid M, Kunhikatta V, Poojari PG, Bakkannavar SM, Balakrishnan JM, Thunga G. The significance of APACHE II as a predictor of mortality in paraquat poisoning: A systematic review and meta-analysis. J Forensic Leg Med. 2023 Jul;97:102548.
20. Khamdamov B. Z., Akhmedov R. M., Khamdamov A. B. The use of laser photodynamic therapy in the prevention of purulent-necrotic complications after high amputations of the lower limbs at the level of the lower leg in patients with diabetes mellitus. Scopus Preview. International journal of Pharmaceutical Research. Volume 11, Issue 3, July-Sept, 2019
21. Khamdamov B. Z., Nuraliev N.A. Pathogenetic approach in complex treatment of diabetic foot syndrome with critical lower limb ischemia. American Journal of Medicine and Medical Sciences, 2020 10 (1) 17-24 DOI: 10.5923/j.20201001.05.
22. Khamdamov B.Z. Indicators of immunocytocine status in purulent-necrotic lesions of the lower extremities in patients with diabetes mellitus. American Journal of Medicine and Medical Sciences, 2020 10 (7): 473-478 DOI: 10.5923/j.20201001.08
23. Khamdamov, B., & Dekhkonov, A. (2022). Clinical and laboratory parameters of the wound process complicated by the systemic inflammatory response syndrome in patients with diabetes mellitus. Journal of education and scientific medicine, 2(3), 25-29. Retrieved from <https://journals.tma.uz/index.php/jesm/article/view/349>
24. Khamroev, U., & Khamdamov, B. (2022). Features of changes in endothelial system parameters in patients with diffuse toxic goiter. Journal of education and scientific medicine, 2(3), 62-67. Retrieved from <https://journals.tma.uz/index.php/jesm/article/view/358>
25. Khamdamova M.T. Age-related and individual variability of the shape and size of the uterus according to morphological and ultrasound studies // News of dermatovenereology and reproductive health. - Tashkent, 2020. - No. 1-2 (88-80). - P.49-52.
26. Khamdamova M. T. Anthropometric characteristics of the physical status of women in the first and second period of middle age // A new day in medicine. Bukhara, 2020. - № 1 (29). - C.98-100.

27. Khamdamova M.T. Age-related and individual variability of the shape and size of the uterus according to morphological and ultrasound studies // News of dermatovenereology and reproductive health. - Tashkent, 2020. - No. 1-2 (88-80). - P.49-52.
28. Khamdamova M.T. Ultrasound features of three-dimensional echography in assessing the condition of the endometrium and uterine cavity in women of the first period of middle age using intrauterine contraceptives // Biology va tibbyot muammolari. - Samarkand, 2020. - No. 2 (118). - P.127-131.
29. Khamdamova M. T. Ultrasound assessment of changes in the endometrium of the uterus in women of the first and second period of middle age when using intrauterine and oral contraceptives // Biomedicine va amaliyot journals. – Tashkent, 2020. - No. 2. - Part 8.- P.79-85.
30. Khamdamova M.T. Features of ultrasound parameters of the uterus in women of the first and second period of middle age using injection contraceptives // New day in medicine. Bukhara, 2020. - No. 2/1 (29/1). - P.154-156.
31. Khamdamova M.T. Features of ultrasound images of the uterus and ovaries in women of the second period of middle age using combined oral contraceptives // New day in medicine. Bukhara, 2020. - No. 2 (30). - P. 258-261.
32. Khamdamova M.T. Individual variability of the uterus and ovaries in women who use and do not use various types of contraceptives // New day in medicine. Bukhara, 2020. - No. 3 (31). - pp. 519-526.
33. Khamdamova M. T. Echographic features variability in the size and shape of the uterus and ovaries in women of the second period of adulthood using various contraceptives // Asian Journal of Multidimensional Research - 2020. – N9 (5). - P.259-263.
34. Khamdamova M. T. Somatometric characteristics of women of the first and second period of adulthood using different contraceptives with different body types // The american journal of medical sciences and pharmaceutical research - 2020. – N8 (2). - P.69-76.
35. Khamdamova M.T., Zhaloldinova M.M., Khamdamov I.B. The state of nitric oxide in blood serum in patients with cutaneous leishmaniasis // New day in medicine. Bukhara, 2023. - No. 5 (55). - P. 638-643.
36. Khamdamova M.T., Zhaloldinova M.M., Khamdamov I.B. The value of ceruloplasmin and copper in blood serum in women wearing copper-containing intrauterine device // New day in medicine. Bukhara, 2023. - No. 6 (56). - pp. 2-7.
37. Khamdamova M. T. Bleeding when wearing intrauterine contraceptives and their relationship with the nitric oxide system // American journal of pediatric

medicine and health sciences Volume 01, Issue 07, 2023 ISSN (E): 2993-2149.
P.58-62

38. Khamdamova M. T. The state of local immunity in background diseases of the cervix // Eurasian journal of medical and natural sciences Innovative Academy Research Support Center. Volume 3 Issue 1, January 2023 ISSN 2181-287X P.171-175.

39. Khamdamova M.T., Khasanova M.T. Various mechanisms of pathogenesis of endometrial hyperplasia in postmenopausal women (literature review) // New day in medicine. Bukhara. 2023. - No. 8 (58). - P. 103-107.

40. Khamdamova M.T. Reproductive Health of Women Using Copper-Containing Intrauterine Contraception // Eurasian Medical Research Periodical Volume 28 January 2024, ISSN: 2795-7624 .www.geniusjournals.org P. 39-45.

41. Khamdamov I.B. Advantages Of Laparoscopic Hernioplasty in Obesity Women of Fertile Age // Eurasian Medical Research Periodical Volume 28 January 2024, ISSN: 2795-7624 .www.geniusjournals.org P. 33-38